

integers) measures the uniform-flux strength, $\Delta\phi$ measures the staggered-flux strength, and both are in units of $\phi_0/2\pi$ ($\phi_0 = hc/e$ is the flux quantum). w_i is a random potential uniformly distributed between $[-W/2, W/2]$.

The gauge selection for this staggered-modulated magnetic field is shown in Fig. 1 and the corresponding periodical boundary condition is adopted. In the absence of disorder ($W = 0$), the size of the chosen magnetic unit cell is $2N \times 2$ for $\phi = \frac{p}{2N} \times 2\pi$. Given N , the variation of $\Delta\phi$ does not change the periodicity of the system. In the presence of disorder ($W \neq 0$), the sample considered has the $2N \times 2N$ unit cell for $\phi = \frac{p}{2N} \times 2\pi$ and the periodical boundary condition is also adopted.

After the numerical diagonalization of the Hamiltonian [Eq. (1)], the total Hall conductance for the system can be calculated through the standard Kubo formula:

$$\sigma_H(E) = \frac{ie^2}{A_0\hbar} \sum_{\varepsilon_m < E} \sum_{\varepsilon_n > E} \frac{\langle m|v_x|n\rangle\langle n|v_y|m\rangle - \langle m|v_y|n\rangle\langle n|v_x|m\rangle}{(\varepsilon_m - \varepsilon_n)^2} \quad (2)$$

where A_0 is the area of the system, E is the Fermi energy, ε_m and ε_n are the corresponding eigenvalues of the eigenstates $|m\rangle$ and $|n\rangle$. The velocity operator is defined as $v_\tau = i \sum_i (c_{i+\tau}^\dagger c_i e^{i\phi_{i+\tau,i}} - c_i^\dagger c_{i+\tau} e^{-i\phi_{i+\tau,i}})$ with $\tau = \hat{x}$ or $\hat{y}$. And we can rewrite σ_H as $\sigma_H(E) = e^2/h \sum_{\varepsilon_m < E} C_m$, where C_m is the Chern number [2, 3] of the m -th subband.




FIG. 2: (color online). The case with $p = 1, N = 4$ and $W = 0$. (a) The Hall conductance σ_H versus the Fermi energy E for various $\Delta\phi$'s. (b)-(e) The DOS of the cases in (a). The Chern numbers of subbands are shown.

Absence of disorder. (a) An example with $p = 1$ and $N = 4$.—An overall picture for the Hall conductance calculated by Eq. (2) is shown in Fig. 2(a) with the uniform-flux strength $\phi = \frac{1}{8} \times 2\pi$ and various staggered-flux strengths $\Delta\phi = 0, 0.1\pi, 0.14\pi, 0.15\pi, 0.2\pi$ (only $E \leq 0$ part is shown here). For the uniform case ($\Delta\phi = 0$), three well-defined IQHE plateaus at $\sigma_H = le^2/h$ ($l = 1, 2, 3$) are clearly shown, corresponding to four Landau subbands [see the corresponding DOS in Fig. 2(b)] centered at the jumps of the Hall conductance at $E \leq 0$. With $\Delta\phi$ increasing from 0 to 0.2π , one sees a systematic evolution

of these IQHE plateaus. At $\Delta\phi = 0.1\pi$, the $l = 3$ IQHE plateau narrows from both sides, and the $l = 2$ IQHE plateau narrows from the right side. At $\Delta\phi = 0.2\pi$, the $l = 3$ IQHE plateau disappears and is replaced by a $l = -1$ IQHE plateau. $\Delta\phi = 0.15\pi$ is close to the critical point where the transition from the $l = 3$ IQHE plateau to the $l = -1$ IQHE plateau takes place. All the values of the Hall conductances in Fig. 2(a) fall back to zero when the Fermi energy approaches $E = 0$, due to the particle-hole symmetry.

Such a behavior can be explained through the evolution of the density of states (DOS) depicted in Fig. 2(b)-(e). In Fig. 2(b),(c) ($\Delta\phi = 0, 0.1\pi$), there are four separated subbands (each totally-filled subband contributes $\frac{1}{8}$ to the filling factor) at $E \leq 0$ with the corresponding well-defined four sequential Chern integers $\{+1, +1, +1, -3\}$. In Fig. 2(d) ($\Delta\phi = 0.15\pi$), the upper two subbands merge together and form a pseudogap which makes the corresponding quantization of Hall conductance not well defined. In Fig. 2(e) ($\Delta\phi = 0.2\pi$), the upper two subbands separate again and the sequential Chern integers are redistributed as $\{+1, +1, -3, +1\}$. We can conclude that at the critical filling $\nu = \frac{3}{8}$, there is a direct transition of the Hall conductance from the $l = 3$ IQHE plateau to the $l = -1$ IQHE plateau when $\Delta\phi$ is increased from 0 up to a critical value $\Delta\phi_c$.

(b) *Redistribution behavior of Chern numbers.*—At $p = 1$, we have numerically confirmed that the Hall conductances at weaker flux strengths with $N = 7-36$ all exhibit the similar features as those in Fig. 2(a). Thus in general, given $\phi = \frac{1}{2N} \times 2\pi$, the N sequential Chern integers $\{+1, \dots, +1, +1, -N+1\}$ will be redistributed as $\{+1, \dots, +1, -N+1, +1\}$ (one can also say that there is a transfer of integer N between the last two Chern numbers) when $\Delta\phi$ is increased from 0 up to a critical value $\Delta\phi_c$. Namely, there is a direct transition of the Hall conductance from the $l = N-1$ IQHE plateau to the $l = -1$ IQHE plateau) at the critical filling $\nu_c = \frac{N-1}{2N}$. This is quite different from the global phase diagram proposed by Kivelson, Lee and Zhang [16] for IQHE which predicts that the only allowed transitions are the nearest-neighbor plateau-plateau transitions ($l \rightarrow l \pm 1$), and is also different from the disorder-induced higher IQHE plateau to insulator transition ($l \rightarrow 0$) [6].

In cases with $p > 1$, based on a Diophantine equation [2], a pattern of subband structure has been found by Yang and Bhatt [7]: the subbands away from the band center form groups with p subbands each, and the total Chern number of each group is $+1$, while the subbands near the band center form a special group with $\text{mod}(N, p)$ subbands.

At $p = 3$ and various N 's, through numerical calculations, we find a more complicated redistribution behavior of Chern numbers (Table I). For instance, we take the case $N = 8$, when $\Delta\phi$ varies from 0 to 0.2π , the eight sequential Chern numbers

TABLE I: Redistribution behavior of Chern numbers (of the $E \leq 0$ subbands) for $p = 3, 5, 7, 9$ and various N 's. Only the last two or three groups are shown for $N \geq 13$.

p	N	$\Delta\phi$	Chern numbers of Landau subbands
3	8	0	$(-5, 11, -5)(-5, 11, -5)(-5, 3)$
		0.03π	$(-5, 11, -5)(-5, 3, 3)(-5, 3)$
		0.1π	$(3, 3, -5)(-5, 3, 3)(-5, 3)$
		0.2π	$(3, 3, -5)(-5, 3, 3)(3, -5)$
3	10	0	$(7, -13, 7)(7, -13, 7)(7, -13, 7)(-3)$
		0.2π	$(-3, -3, 7)(7, -3, -3)(-3, -3, 7)(-3)$
3	11	0	$(-7, 15, -7)(-7, 15, -7)(-7, 15, -7)(-7, 4)$
		0.15π	$(-7, 4, 4)(4, 4, -7)(-7, 4, 4)(4, -7)$
3	13	0	$\dots (9, -17, 9)(9, -17, 9)(9, -17, 9)(-4)$
		0.1π	$\dots (-4, -4, 9)(9, -4, -4)(-4, -4, 9)(-4)$
3	16	0	$\dots (11, -21, 11)(11, -21, 11)(-5)$
		0.1π	$\dots (11, -5, -5)(-5, -5, 11)(-5)$
3	17	0	$\dots (-11, 23, -11)(-11, 23, -11)(-11, 6)$
		0.08π	$\dots (6, 6, -11)(-11, 6, 6)(6, -11)$
3	32	0	$\dots (-21, 43, -21)(-21, 43, -21)(-21, 11)$
		0.05π	$\dots (11, 11, -21)(-21, 11, 11)(11, -21)$
5	12	0	$(5, 5, -19, 5, 5)(5, 5, -19, 5, 5)(5, -7)$
		0.2π	$(5, 5, -7, -7, 5)(5, -7, -7, 5, 5)(-7, 5)$
5	13	0	$\dots (-5, -5, 21, -5, -5)(-5, -5, 8)$
		0.2π	$\dots (-5, -5, 8, 8, -5)(-5, 8, -5)$
5	14	0	$\dots (-11, 17, -11, 17, -11)(-11, 17, -11, 3)$
		0.2π	$\dots (-11, 3, 3, 3, 3)(3, 3, -11, 3)$
5	17	0	$\dots (7, 7, -27, 7, 7)(7, 7, -27, 7, 7)(7, -10)$
		0.08π	$\dots (7, 7, -10, -10, 7)(7, -10, -10, 7, 7)(-10, 7)$
5	33	0	$\dots (-13, -13, 53, -13, -13)(-13, -13, 20)$
		0.04π	$\dots (-13, -13, +20, +20, -13)(-13, 20, -13)$
7	16	0	$\dots (-9, 23, -9, -9, -9, 23, -9)(-9, 7)$
		0.08π	$\dots (-9, 7, 7, -9, -9, 7, 7)(7, -9)$
7	23	0	$\dots (-13, 33, -13, -13, -13, 33, -13)(-13, 10)$
		0.06π	$\dots (-13, 10, 10, -13, -13, 10, 10)(10, -13)$
7	30	0	$\dots (-17, 43, -17, -17, -17, 43, -17)(-17, 13)$
		0.04π	$\dots (-17, 13, 13, -17, -17, 13, 13)(13, -17)$
7	31	0	$\dots (9, 9, 9, -53, 9, 9, 9)(9, 9, -22)$
		0.04π	$\dots (9, 9, -22, -22, 9, 9, 9)(9, -22, 9)$
9	20	0	$\dots (9, 9, -31, 9, 9, 9, -31, 9, 9)(9, -11)$
		0.06π	$\dots (9, -11, -11, 9, 9, -11, -11, 9, 9)(-11, 9)$
9	22	0	$\dots (5, 5, 5, 5, -39, 5, 5, 5, 5)(5, 5, 5, -17)$
		0.06π	$\dots (5, 5, 5, -17, -17, 5, 5, 5, 5)(5, 5, -17, 5)$
9	29	0	$(13, 13, -45, 13, 13, 13, -45, 13, 13)(13, -16)$
		0.04π	$(13, -16, -16, 13, 13, -16, -16, 13, 13)(-16, 13)$
9	38	0	$(17, 17, -59, 17, 17, 17, -59, 17, 17)(17, -21)$
		0.03π	$(17, -21, -21, 17, 17, -21, -21, 17, 17)(-21, 17)$

$\{(-5, +11, -5), (-5, +11, -5), (-5, +3)\}$ (each pair of parentheses represents a group according to the grouping scheme by Yang and Bhatt [7]) are redistributed as $\{(+3, +3, -5), (-5, +3, +3), (+3, -5)\}$. We note that, in each group, one Chern number is subtracted by 8 and another neighboring Chern number is added by 8. Hence, the total quantized Hall conductance exhibits a direct transition by $\pm 8e^2/h$ at the critical fillings $\nu_c = \frac{1}{16}, \frac{5}{16}$ and $\frac{7}{16}$. At $N = 10$, there is only one Chern number in the last group which remains unchanged when $\Delta\phi$ varies, while the other three groups change similarly as the case $N = 8$. We have numerically confirmed that for $p = 5, 7$ and 9 (Table I), there are also redistribution behaviors similar to $p = 3$.

In general, at $p \geq 3$, there is a redistribution behavior of Chern numbers among each group [except the last group when $\text{mod}(N, p) = 1$ and there is only one Chern number within it]: a transfer of integer N between two neighboring Chern numbers, and the redistributed Chern numbers of neighboring groups away from the band center are antisymmetric in sequence. Hence, the total quantized Hall conductance will exhibit a direct transition by $\pm Ne^2/h$ at critical fillings.

(c) *Scaling properties.*—Both the uniform-flux strength $\phi = \frac{p}{2N} \times 2\pi$ at $N = 4-38$ and the staggered-flux strength in the previous calculations are relatively high fields. In order to meet the requirements of possible experimental realization, we will reduce the flux strength further. The scaling behavior of $\Delta\phi_c$ with ϕ is shown in Fig. 3. For $p \geq 3$, we concentrate on the cases where there are two Chern numbers in the last group and consider the critical filling $\nu = \frac{N-1}{2N}$. For a fixed p , $\Delta\phi_c$ scales with ϕ almost linearly. The slopes of the lines with different p 's differ from one another. With ϕ decreasing, all lines approach the origin point, and therefore $\Delta\phi_c$ approaches zero at the weak-field limit.




FIG. 3: (color online). $\Delta\phi_c$ versus ϕ for $p = 1, 3, 5, 7, 9$ and various N 's (at the critical filling $\nu = \frac{N-1}{2N}$)

Presence of disorder.—In realistic solids, impurity or phonon-induced disorder always has effect on the

magneto-transport properties. We now consider $W = 1$ to see the influence of weak disorder on the critical transition behavior.




FIG. 4: (color online). The case with $p = 1$, $N = 4$ and $W = 1$ (1000 random-potential configurations of the lattice size 8×8). σ_H versus E for various $\Delta\phi$'s. The inset shows the evolution of σ_H versus $\Delta\phi$ at $\nu = \frac{3}{8}$.

As for $p = 1$ and $N = 4$, we can see from Fig. 4 that the presence of weak disorder does not smear out the transition of the Hall conductance from the $l = 3$ IQHE plateau to the $l = -1$ IQHE plateau near the critical filling $\nu = \frac{3}{8}$.

An experimental probe of the SFP order.—As one application of our theoretical concern, we would take the staggered-flux part $\pm\Delta\phi$ as the possible internal orbital-antiferromagnetic structure of the SFP order in some strongly-correlated electron systems [10, 12], and the uniform-flux part ϕ as an external experimental probe. In Fig. 5, at the electron filling $\nu = \frac{15}{32}$, we fix the value of $\Delta\phi$ and do such a hypothetical experiment by measuring the Hall conductance which varies with ϕ . The two curves of the Hall conductances with $\Delta\phi = 0$ and $\Delta\phi = 0.1\pi$ exhibit large deviations at many values of N/p . Therefore, such an experiment could tell us the characteristic differences between the states with and without the SFP order.

Summary and discussion.—As for the tight-binding electrons on a square lattice under a staggered-modulated magnetic field with the strength $\phi \pm \Delta\phi$, a new peculiar behavior of the Hall conductances has been found: (a) if $\phi = \frac{p}{2N} \times 2\pi$ (p and $2N$ are coprime integers) is fixed and $\Delta\phi$ is increased from 0 up to a critical value $\Delta\phi_c$, the Chern numbers of Landau subbands will be redistributed between neighboring pairs; (b) and hence the total quantized Hall conductance exhibits a direct transition by $\pm Ne^2/h$ at one or several critical fillings; (c) the neighboring two subbands merge together and form a pseudogap when $\Delta\phi = \Delta\phi_c$; (d) at a fixed p , $\Delta\phi_c$ scales with ϕ almost linearly; (e) weak disorder does not smear out the peculiar transition of the Hall conductance.

To observe this effect under an accessible uniform mag-




FIG. 5: (color online). At $\nu = \frac{15}{32}$, σ_H versus π/ϕ for $\Delta\phi = 0$ and 0.1π . N ranges from 8 to 64.

netic field with $N \sim 1000$, we estimate that both the strength of disorder W and the temperature $k_B T$ should not exceed $\frac{4t}{N} \sim \frac{4t}{1000}$ (where $4t$ is half of the original band width). Taking $t = 0.01$ eV, the temperature T should be lower than 0.5 K.

As for high- T_c cuprates, the strength of the internal staggered magnetic field is estimated to be about 0.1 T [17]. One may suspect that: will an internal staggered magnetic field survive under a stronger external uniform one? We believe it does, basing on the numerical results: an external uniform magnetic field will suppress the d -wave superconducting order, while the SFP order (or equivalently, the DDW order) can exist or even be enhanced in the vortex cores [18, 19].

This work was supported by the State Key Program for Basic Researches of China under Grant No. 2006CB921802 and the Research Grants Council of Hong Kong.

-
- [1] The Quantum Hall Effect, 2nd Ed., edited by R. E. Prange and S. M. Girvin (Springer-Verlag, New York, 1990).
 - [2] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, Phys. Rev. Lett. **49**, 405 (1982).
 - [3] D. P. Arovas, R. N. Bhatt, F. D. M. Haldane, P. B. Littlewood, and R. Rammal, Phys. Rev. Lett. **60**, 619 (1988).
 - [4] D. Z. Liu, X. C. Xie, and Q. Niu, Phys. Rev. Lett. **76**, 975 (1996).
 - [5] K. Yang and R. N. Bhatt, Phys. Rev. Lett. **76**, 1316 (1996).
 - [6] D. N. Sheng and Z. Y. Weng, Phys. Rev. Lett. **78**, 318 (1997); Phys. Rev. Lett. **80**, 580 (1998).
 - [7] K. Yang and R. N. Bhatt, Phys. Rev. B **59**, 8144 (1999).
 - [8] S. V. Kravchenko, W. Mason, J. E. Furneaux, and V. M. Pudalov, Phys. Rev. Lett. **75**, 910 (1995).
 - [9] S. H. Song, D. Shahar, D. C. Tsui, Y. H. Xie, and D. Monroe, Phys. Rev. Lett. **78**, 2200 (1997); M. Hilke, D. Shahar, S. H. Song, D. C. Tsui, and Y. H. Xie, Phys. Rev. B **62**, 6940 (2000).

- [10] S. Chakravarty, R. B. Laughlin, D. K. Morr, and C. Nayak, Phys. Rev. B **63**, 094503 (2001).
- [11] T. Timusk and B. Statt, Rep. Prog. Phys. **62**, 61 (1999).
- [12] P. Chandra, P. Coleman, J. A. Mydosh, and V. Tripathi, Nature (London) **417**, 831 (2002).
- [13] Y. Iye, E. Kuramochi, M. Hara, A. Endo, and S. Katsumoto, Phys. Rev. B **70**, 144524 (2004).
- [14] Q. W. Shi and K. Y. Szeto, Phys. Rev. B **56**, 9251 (1997).
- [15] G. Y. Oh, Phys. Rev. B **60**, 1939 (1999).
- [16] S. Kivelson, D. H. Lee, and S. C. Zhang, Phys. Rev. B **46**, 2223 (1992).
- [17] T. C. Hsu, J. B. Marston, and I. Affleck, Phys. Rev. B **43**, 2866 (1991).
- [18] Q. H. Wang, J. H. Han, and D. H. Lee, Phys. Rev. Lett. **87**, 167004 (2001).
- [19] J. An and C. D. Gong, Phys. Rev. B **70**, 012505 (2004).